

Josephson Junctions with Minimal Length

T. L. Boyadjiev

*Faculty of Mathematics and Computer Science,
Sofia University 'St. Kliment Ohridski', Bulgaria
e-mail: todorlb@fmi-uni.sofia.bg*

M. D. Todorov

*Faculty of Applied Mathematics and Computer Science,
Technical University of Sofia, Bulgaria
e-mail: mtod@vmei.acad.bg*

PACS: 85.25.Cp, 85.25.Am, 02.60.Lj

Abstract

The minimal length of “one-dimensional” Josephson junctions, in which the specific bound states of the magnetic flux retain their stability is discussed numerically. Thereby, we consider as “long” every Josephson junction, in which there exists at least one nontrivial stable distribution of the magnetic flux for fixed values of all the physical and the geometrical parameters.

Our results can be applied for optimization of the sizes of devices containing JJs for different operating conditions.

1 Formulation of the Problem

It is known that the stationary distributions (bound states) of the magnetic flux $\varphi(x)$ in “long” (one-dimensional) Josephson junctions (JJs) can be modeled as solutions of the nonlinear boundary value problem (BVP):

$$-\varphi_{xx} + j_D(x) \sin \varphi + \gamma = 0, \quad \varphi_x(\pm R) = h_B, \quad (1)$$

where γ is the bias current, and h_B is the external (boundary) magnetic field. We note that the kind of boundary conditions, either in presence or in absence of the current γ in the right-hand side of eqn. (1), is determined by the geometry of the junction. Here we consider simple JJs with overlap geometry [1], in which the current γ can be approximately considered as a constant. The generalization of our results in the case of another geometry, for example in-line geometry, does not require big efforts.

We suppose that the given function $0 \leq j_D(x) \leq 1$ describes the variations of the Josephson current amplitude, caused by the possible local inhomogeneities of the dielectric layer thickness [2]. When $j_D(x) \equiv 1$ the junction is homogeneous (see the complete analytical results in recent paper [3]).

The single inhomogeneity is characterized by its respective position x_0 and width μ , so the amplitude $j_D = j_D(x, x_0, \mu)$. The frequently used expression $j_D(x) = 1 - \mu\delta(x - x_0)$ (here $\delta(x)$ is Dirac's function) is useful in theoretical examination (see the basic papers [4] and [5]). In this paper we use an isosceles trapezium with base μ (Fig.1) as a more suitable physical model of inhomogeneity [6, 7].

Amplitude of the Josephson current




Figure 1

The stability of some concrete solution $\varphi(x)$ of the BVP (1) is determined by the sign of the minimal eigenvalue λ_{min} of the corresponding Sturm-Liouville problem (SLP):

$$-\psi_{xx} + q(x)\psi = \lambda\psi, \quad \psi_x(\pm R) = 0, \quad (2)$$

where $q(x) = j_D(x) \cos \varphi(x)$ is the potential of SLP, originated from the mentioned solution $\varphi(x)$.

Each of the different eigenvalues $\lambda_{min} \equiv \lambda_0 < \lambda_1 < \lambda_2 < \dots < \lambda_n < \dots$, conforms to an unique eigenfunction $\psi_n(x)$, $n = 0, 1, 2, \dots$, which satisfies some additional (norm) condition, for example:


$$\int_{-R}^R \psi^2(x) dx = 1. \quad (3)$$

If the minimal eigenvalue $\lambda_{min} > 0$, then the respective solution $\varphi(x)$ of BVP (1) is stable with respect to small time-space perturbations. If the minimal eigenvalue $\lambda_{min} < 0$, then this solution is unstable. The value $\lambda_{min} = 0$ corresponds to a bifurcation point, in which the stable solutions of eqn. (1) go to unstable ones and vice versa (for details see [5]).

Apart from the space coordinate x , the virtual solutions of the nonlinear BVP (1) depend also on the physical parameters h_B , γ and the “technological” ones μ , $\Delta = 2R$ (length of the junction), and x_0 , i.e., $\varphi = \varphi(x, p)$, where we simply substitute $p \equiv \{h_B, \gamma, \mu, x_0, \Delta\}$. The varying of each of those parameters causes a variation of the distribution $\varphi(x, p)$ and, therefore, subsequent variations of the potential $q(x, p)$, the eigenvalues $\lambda(p)$, and the respective eigenfunctions $\psi(x, p)$. Thus, we can conclude that every concrete solution of BVP (1) has an area, where this solution remains stable with regard to the variations of the parameters p . In the parametric space the equation:

$$\lambda_{min}(p) = 0 \quad (4)$$

determines a hypersurface, whose points appear to be bifurcation points corresponding to the solution under consideration. The intersections of the bifurcation hypersurface (4), when there are chosen pairs of the parameters p for fixed values of others, are called bifurcation curves and the respective values of the parameters - bifurcation (critical) parameters. From a physical viewpoint most interesting seem to be the bifurcation curves of the kind “current - magnetic flux” $\lambda_{min}(h_B, \gamma) = 0$, when the geometrical parameters μ , R , and x_0 are fixed. These curves could be obtained experimentally [8, 9, 10].



From a technological point of view, however, it is worth investigating the bifurcation curves as functions with respect to at least one of the geometrical parameters, for example:

$$\lambda_{min}(h_B, \Delta) = 0, \lambda_{min}(\gamma, \Delta) = 0, \lambda_{min}(\mu, \Delta) = 0, \dots$$

Such kind of problems can be connected with the optimization of sizes of devices, containing JJs.

Most of our results below are related to the typical solution of BVP (1), physically existing both in homogeneous and inhomogeneous JJs: so called single fluxon. Let us note that in the infinite homogeneous or inhomogeneous JJ with one δ -shaped microinhomogeneity at the point $x_0 = 0$, the single fluxon/antifluxon is represented by the exact solution [5], [1]:

$$\varphi(x) = 4 \arctan \exp(\pm x). \quad (5)$$

More complicated (multi-fluxon) solutions there exist in inhomogeneous JJs for big enough values of h_B .

The numerically obtained basic functions: the magnetic flux $\varphi(x)$, the magnetic field $\varphi_x(x)$, and the first eigenfunction $\psi_0(x)$ of such solution for $\Delta = 6.3$, $\mu = 1$, $h_B = 0$, and $\gamma = 0$ are represented at Fig. 2.

Fig. 3 represents the typical relationships $\lambda_{min}(\Delta)$, corresponding to the single fluxons in inhomogeneous junctions with one resistive inhomogeneity placed at the point $x_0 = 0$ when $\mu = 0.5$ and $\mu = 1$, respectively. The points B_0 and B_1 appear to be bifurcation points for these solutions, i.e., the zeroes of the corresponding relationships $\lambda_{min}(\Delta)$. According to the above mentioned reasonings, these points determine the minimal length of the JJ, for which the single fluxon is still stable for given values of the other parameters.




Figure 3

Usually, from physical point of view, the concrete JJ is “long” or “one-dimensional” if the Josephson penetration depth λ_J satisfies the condition $\lambda_J \gg 1$. However, such definition is not perfect enough mathematically. On the other hand, it seems that the calculation of the minimal geometrical sizes and, especially, the minimal length Δ of homogeneous or inhomogeneous JJs, corresponding to a concrete nontrivial distribution of the magnetic flux, is an important practical problem.

The trivial method, obviously, is through the point by point construction of the curves of type $\lambda_{min}(\Delta)$ using BVPs (1)-(3). However, it is quite natural to put the question how to calculate directly the bifurcation curves. A general approach of study for the formulated problem is proposed in the papers [11, 12].

We consider the system of the equations (1), (2), and (3) as a closed nonlinear eigenvalue problem with respect to the functions $\varphi(x)$, $\psi(x)$ and one of the parameters p , for example, h_B or γ in [12], or R in this case, while the other parameters, and the eigenvalue λ as well, are given. Fixing $\lambda > 0$ to be small enough (for example $\lambda = 0.001 \div 0.01$), we guarantee a stable solution to be obtained. We note, that the derivative $\partial\lambda/\partial p_0 \rightarrow \infty$ when p_0 approaches its critical (bifurcation) value, so, the solutions of the above system with a priori prescribed accuracy belong to the small vicinity of the sought bifurcation curve.

The generalized Continuous Analog of Newton’s Method [13] for solving the mentioned system is used (see details in our recent paper [7]).

We have to note that for some initial approximations the method can lead to an unstable solution, as well. In this case, the given λ corresponds to one of the higher eigenvalues, not to the minimal one. The simplest method to avoid this difficulty uses the oscillation theorem [14] for eigenfunctions: *If the obtained function $\psi(x)$ is the minimal eigenfunction, then it does not have a zero on the interval Δ .*

2 Results and Discussion

Fig. 4 shows the convergence of the iteration process. The upper curve, marked by “ ∇ ”, presents the initial distribution of the magnetic field $\varphi_x(x)$ alongside the junction. This distribution is chosen to be a solution of BVP (1), for $\Delta = 10$, $\mu = 1$, $h_B = 0$, and $\gamma = 0$. At the same figure the lower curve, marked by “o”, presents the calculated distribution of the magnetic field corresponding to the minimal eigenvalue $\lambda_{min} = 0.01$. Hence, this is the sought bifurcation distribution and, in the framework of this model, the value of the spectral parameter $\Delta_{min} \approx 4.24$ is the minimal length of the junction, providing a stable single fluxon. In this sense, junctions, whose length lies below the critical value Δ_{min} of Δ , have to be considered as “short” for distributions $\varphi(x)$ of the kind single fluxon. However, for such a short length Δ there are a unique stable and unstable distributions of the magnetic flux in the junction - Meissner’s solutions: the trivial solutions $\varphi(x) = 0, 2\pi, \dots$ (stable) and $\varphi(x) = \pi, 3\pi, \dots$

(unstable) for $h_B = 0$ and $\gamma = 0$.




Figure 4

The influence of the parameters h_B , γ , μ , and x_0 on the minimal length Δ_{min} of the JJs will be considered below.

If the current $\gamma \neq 0$ and the values of all other parameters are fixed, then the stability of the fluxon requires the junction to be longer (see Fig. 5). This fact corresponds to the classical conclusion [4] that the existence of quantity γ violates the stability of the bound states in JJs. The “force” γ displaces the maximum of $\varphi_x(x)$ from $x_0 = 0$ to the left or right depending on its sign; the full magnetic flux $\Delta\varphi = \varphi(R) - \varphi(-R)$ increases, so that the minimal junction’s length Δ_{min} , necessary to accommodate such a flux, increases as well.




Figure 5




Figure 6

The numerically obtained influence of the current γ upon the bifurcation length Δ_{min} is plotted on Fig.6 for two values of the size μ of the inhomogeneity: $\mu = 0.5$ and $\mu = 1$. The curves increase very abruptly and when γ approaches some critical value, the derivative $\partial\Delta_{min}/\partial\gamma \rightarrow \infty$. This behaviour means that beyond the critical value of γ , JJs which correspond to the single fluxon do not exist.




Figure 7

On the contrary, if the boundary magnetic field $h_B \neq 0$ and varies within some fixed domain, then the stability of the fluxon is improved (see Fig. 7). Well noticeable is the stabilizing influence of the boundary magnetic field h_B upon the critical length of the junction, i.e., the increase of the magnitude of h_B decreases the minimal length of the junction Δ_{min} . Thus, the calculated minimal length of the JJ corresponding to the single fluxon ($h_B = 0$, $\gamma = 0$) is $\Delta_{min} \approx 4.24$, while the same quantity when $h_B = 1$ and $\gamma = 0$ is $\Delta_{min} \approx 2.7$.




Figure 8

Fig.8 represents the distributions of magnetic field $\varphi_x(x)$ alongside the junction in absence of current ($\gamma = 0$) depending on its boundary values $h_B = 0$ and $h_B = 1$, respectively. It is clear that the two solutions could be well approximated by means of two-degree polynomials. Furthermore, they seem to be geometrically similar. Namely, the curve $\varphi_x(x)$ corresponding to a magnetic field $h_B > 0$ may be considered as obtained from the other curve by a homothety alongside x axis with respect to the pole $x = 0$ and a translation alongside the axis φ_x .




Figure 9

At the next Fig.9, the dependence of the minimal length Δ_{min} on the boundary magnetic field h_B for $\gamma = 0$ in the case of a homogeneous JJ (the amplitude $j_D(x) \equiv 1$) is presented. Even though this dependence is qualitatively similar with the analogous ones at Fig.7, it differs from the big slope of Δ_{min} for small h_B , so that $\Delta_{min}(h_B) \rightarrow \infty$ when $h_B \rightarrow 0$.

Such behaviour of the curve $\Delta_{min}(h_B)$ can be qualitatively explained by means of the formula [5]:

$$\lambda_0 = \lambda_{min} = \frac{\mu}{2} \left[\sqrt{1 + \frac{\mu^2}{16}} - \frac{\mu}{4} \right], \quad (6)$$

which is exact for the single fluxon (5) in an infinite JJ with one resistive δ -shaped inhomogeneity at the center $x = 0$. Hence, for $\mu = 0$ we obtain $\lambda_{min} = 0$, so that for $h_B = 0$ and $\gamma = 0$ the single

fluxon in homogeneous JJ has to be considered as a quasi-stable (bifurcation) state of the magnetic flux $\varphi(x)$ [5]. On the contrary, in inhomogeneous JJ with resistive inhomogeneities the single fluxon is still stable ($\lambda_{min} > 0$), which physically is due to the existence of the inhomogeneity.

In the case of an inhomogeneous JJ with finite length $\Delta = 6.3$ the calculated curve $\lambda_{min}(\mu)$ (marked by “□”) for $h_B = 0$ and $\gamma = 0$ is shown at Fig.10. We note that for small enough values of the size μ the dependence is analogous to (6).

Let us introduce the full energy F connected with some solution $\varphi(x)$ of BVP (1) via the formula:

$$F(\mu) = \int_{-R}^R \left[\frac{1}{2} \varphi_x^2 + j_D(x) (1 - \cos \varphi) + \gamma \varphi \right] dx + h_B [\varphi(R) - \varphi(-R)].$$

It is efficient to compare the energy curves $F(\mu)$ (marked by “▽”) for fixed $\Delta = 6.3$ and $F_{min}(\mu)$ (marked by “○”) for a JJ with minimal length (see Fig. 10). The magnitudes of the energy are regarded to the energy ($F = 8$) of the single fluxon (5) in an infinite homogeneous JJ. It is well noticeable that for $\mu > 0$ the inequality $F(\mu) > F_{min}(\mu)$ is satisfied, e.g., the single fluxon in the JJ with minimal length is more stable. The cusp for $\mu = 0$ corresponds to a bifurcation in accordance with formula (6).




Figure 10




Figure 11

At Fig.11 the numerically obtained relationship $\Delta_{min}(\mu)$ is drawn, depending on the length of the trapezium's base μ (width of the inhomogeneity) (see Fig.1). The presented three curves corresponding to different locations of the centre of inhomogeneity x_0 are qualitatively similar. The most important their feature is the existence of minimums, which are placed in the interval $\mu \in (0.9, 1.3)$. Therefore, there exist such values of the quantity μ , which provide a minimal bifurcation length of a JJ from among all the minimal lengths for given other parameters. Let us call them "optimal" lengths. For example, if $x_0 = 0$ ("symmetric" inhomogeneous JJ), the optimal length $\Delta_{opt} \equiv \Delta_{min} \simeq 4.22$ of the JJ corresponds to $\mu \simeq 1.1$, whereas $\Delta_{min} \simeq 4.48$ if the width of the inhomogeneity is $\mu = 0.5$, and $\Delta_{min} \simeq 4.3$ for $\mu = 1.5$.




Figure 12

The bifurcations of the bound states in the circular JJs with respect to the location x_0 of the δ -shaped inhomogeneity are investigated numerically in [15]. Fig.12 shows the curves $\Delta_{min}(x_0)$ describing the behaviour of the minimal length of the linear JJs. When the current $\gamma = 0$ and the width of inhomogeneity $\mu = 1$ is fixed, the curve is symmetric and, apparently, the minimal value of Δ_{min} is reached at the location $x_0 = 0$. When the current becomes nontrivial ($\gamma = 0.1$), the minimum shifts into some other point. For negative values of γ the dependence is the same but mirror reflected with respect to the line $x_0 = 0$.




Figure 13

Finally, the last Fig.13 provides as an illustration a three-soliton bifurcation solution. Such kind of solutions can be obtained when the boundary magnetic field h_B is big enough. Here, the concrete solution is derived for $h_B = 3.2$. At the same graph the initial distribution of the magnetic field $\varphi_x(x)$ as solution of BVP (1), for $\Delta = 8$, $\gamma = 0$, and $\mu = 1$ is included.

Acknowledgements. We thank Prof. I.V. Puzynin (JINR, Dubna, Russia) for useful remarks. TLB remembers with gratitude the elaborate and fruitful discussions with Prof. Yu.S. Gal'pern, Prof. A.T. Filippov, and Prof. A.V. Ustinov in JINR, Dubna, Russia.

References

- [1] Barone A and G. Paterno G 1982 *Physics and Applications of the Jisephson Effect* (John Wiley & Sons, Inc.)
- [2] Anderson P W 1962 *Phys. Rev. Lett.* **9** 309
- [3] Caputo J-G, Flytzanis N, Gaididei Y, Stefanakis N, and Vavalis E (2000) *Superconductor Science and Technology* **13** 423
- [4] McLaughlin D W and Scott A C 1978 *Phys. Rev. A* **18** 1652
- [5] Gal'pern Yu S and Filippov A T 1984 *Sov.JETP* **86**(4) 1527 (in Russian)
- [6] Alexeeva N V and Boyadjiev T L 1997 *Bulg. J. of Physics*(1,2)
- [7] Boyadjiev T L and Todorov M D 2000 *Mathematical Modeling (Russian Academy of Sciences)* **12**(4) 61
- [8] Vystavkin A N, Drachevski Yu F, Kosheletz V P, and Serpuchenko I L 1988 *Fizika nizkih temperatur (Russian Phys. of Low Temper.)* **14**(6) 646 (in Russian)
- [9] Larsen B H, Mygind J, and Ustinov A V 1994 *Phys. Lett. A* **193** 359
- [10] Malomed B A, Ustinov A V 1994 *Phys. Rev. B* **49**(18) 13024
- [11] Filippov A T, Gal'pern Yu S, Boyadjiev T L, and Puzynin I V (1987) *Phys. Lett. A* **120**(1)
- [12] Boyadjiev T L, Pavlov D V, and Puzynin I V 1988 *Comm. of JINR, Dubna* P11-88-409 (in Russian)
- [13] Puzynin I V, Amirkhanov I V, Zemlyanaya E V, Pervushin V N, Puzynina T P, Strizh T A, and Lakhno V D 1999 *Physics of Elementary Particles and Atomic Nuclei*(JINR, Dubna) vol. 30, No 1, p.210 (in Russian), English translation:(American Institute of Physics)p.97
- [14] Levitan B M and Sargsjan I S 1975 *Introduction to Spectral Theory*, Transl. Math. Monographs(RI:AMS, Providence)
- [15] Kaschieva V 1997 *Lecture Notes Comp. Sci.* **1196** (Springer)pp 236–242